


AD-A183 744 THE METABOLISM OF NITROGUANIDINE AND NITROSOGUANIDINE 1/1
BY RAT HEPATIC SUBC. (U) LETTERMAN ARMY INST OF
RESEARCH PRESIDIO OF SAN FRANCISCO CA
UNCLASSIFIED P B SIMBOLI ET AL. JUL 87 LAIR-87-78 F/G 6/1 NL





MICROCOPY RESOLUTION TEST CHART

NATIONAL BUREAU OF STANDARDS 1963-A



(2)

DTIC FILE COPY

DTIC
ELECTE
AUG 1 1 1987
S
CKD
D

Laboratory Note No. 87-70

AD-A183 744

The Metabolism of
Nitroguanidine and Nitrosoguanidine
by
Rat Hepatic Subcellular Fractions

Paul B. Simboli, BS, SGT
Bert Ho, PhD
and
Don W. Korte, Jr., PhD, MAJ MSC

TOXICOLOGY BRANCH
DIVISION OF COMPARATIVE MEDICINE AND TOXICOLOGY

DISTRIBUTION STATEMENT A

Approved for public release
Distribution Unlimited

JULY 1987

Toxicology Series 152

LETTERMAN ARMY INSTITUTE OF RESEARCH
PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

The metabolism of nitroguanidine and nitrosoguanidine by rat hepatic subcellular fractions--Simboli et al.


Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

In conducting the research described in this report, the investigation adhered to the "Guide for the Care and Use of Laboratory Animals," as promulgated by the Committee on Revision of the Guide for Laboratory Animal Facilities and Care, Institute of Laboratory Animal Resources, National Research Council.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)


(Signature and date) 7/1/57

This document has been approved for public release and sale; its distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Laboratory Note No. 87-70	2. GOVT ACCESSION NO. AD-A183744	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) The metabolism of nitroguanidine and nitroso- guanidine by rat hepatic subcellular fractions		5. TYPE OF REPORT & PERIOD COVERED
7. AUTHOR Paul B. Simbol, BS, SGT; Bert Ho, PhD; Don W. Korte, Jr, PhD, MAJ MSC		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Division of Toxicology Letterman Army Institute of Research Presidio of San Francisco, CA 94129-6800		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
12. REPORT DATE July 1987		13. NUMBER OF PAGES 7
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) NITROGUANIDINE, NITROSOGUANIDINE		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The metabolism of nitroguanidine (NG) and nitrosoguanidine (NsG) by rat hepatic subcellular fractions was examined. The microsomal fraction and 10,000 x g supernatant from untreated and phenobarbital-treated animals and the 9,000 x g supernatant from Arochlor-induced rats were used for this study. The <u>in vitro</u> metabolism of substrate was measured by HPLC. The results indicate that neither NG nor NsG are metabolized by the hepatic subcellular fractions.		

ABSTRACT

The metabolism of nitroguanidine (NG) and nitrosoquanidine (NsG) by rat hepatic subcellular fractions was examined. The microsomal fraction and 10,000 x g supernatant from untreated and phenobarbital-treated animals and the 9,000 x g supernatant from Arochlor-induced rats were used for this study. The in vitro metabolism of substrate was measured by HPLC. The results indicate that neither NG nor NsG are metabolized by the hepatic subcellular fractions.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



The Metabolism of Nitroguanidine and Nitrosoguanidine by
Rat Hepatic Subcellular Fractions--Simboli et al

Nitroguanidine, a primary component of US Army triple-base propellants, is now produced in a Government-owned contractor-operated ammunition plant. The US Army Biomedical Research and Development Laboratory (USABRDL), as part of its mission to evaluate the environmental and health hazards of military-unique pollutants generated by US Army munitions manufacturing facilities, conducted a review of the nitroguanidine data base and identified significant gaps in the toxicity data. The Toxicology Branch, LAIR, was tasked by USABRDL to develop a genetic and mammalian toxicity profile for nitroguanidine, related intermediates/by-products of its manufacture, and its environmental degradation products. Nitrosoguanidine is a potential environmental degradation product of nitroguanidine.

The development of a toxicity profile for NG and NsG will aid in understanding the safety or hazards of these compounds. Metabolic fate studies are an integral part of these toxicological tests. To supplement in vivo studies on the fate of nitroguanidine in the rat, this study examines the in vitro metabolism of NG and NsG by hepatic subcellular fractions. Additional impetus to examine the in vitro metabolism of NsG is provided by preliminary reports that NsG was mutagenic in mouse lymphoma assays when activated with Arochlor-induced 9,000 x g supernatant (9K) (1).

The objective of this study is to quantitate the extent of metabolism of NG and NsG by three rat hepatic subcellular fractions: untreated and phenobarbital-treated 10,000 x g supernatant (10K), Arochlor-induced 9,000 x g supernatant, and untreated and phenobarbital-treated microsomal pellet. If the substrates are found to be degraded metabolically, the metabolites will be isolated, identified, and quantified.

MATERIALS AND METHODS

Chemicals

Glucose-6-phosphate, glucose-6-phosphate dehydrogenase, NADP, and Fischer's medium for leukemic cells of mice were

purchased from the Sigma Chemical Co., St. Louis, MO. Nitroguanidine was supplied by the Sunflower Ammunition Plant, Desoto, KS. The US Army Biomedical Research and Development Laboratory (USABRDL, Fort Detrick, MD) provided the NSG. The Arochlor-induced PK supernatant was supplied by Microbiological Associates, Bethesda, MD.

Instrumentation

The HPLC system consisted of two LKB 2150 pumps coupled to an LKB 2152 HPLC controller; 20- μ l injections were made via a Waters Wisp 710B; the detector was a Fritios Spectroflow 773; the integrator was a Shimadzu C-P3A. The HPLC column was a Brownlee C-18 Spheri-5 column (4.6 mm x 250 mm); a mobile phase of 10% MeOH/H₂O was used at a flow rate of 0.7 ml/min; the detector was set at 265 nm (1.000 AUFS). The centrifuges used were a Sorvall GRC-1, a Sorvall RC-5 Superspeed Refrigerated centrifuge, and a Beckman L5-75 Ultracentrifuge. A Eimontec PCU homogenizer was used to homogenize the livers.

Animals

Male albino Sprague-Dawley rats (350-425 g) were obtained from Pantin-Kingman, Fremont, CA. The inducing agent, sodium phenobarbital (PB), was administered ip (80 mg/kg) for four consecutive days prior to sacrifice on day 5. The untreated animals were left untreated during this time period. Animals were fasted for a 24 hour period prior to sacrifice (SOP-OP-STX-88, "Preparation of Hepatic Subcellular Fractions"). The rats were sacrificed by decapitation and the livers were rapidly removed and rinsed in cold 1.15% KCl. The livers were blotted dry, weighed, and placed in a volume of cold 0.1M Tris buffer (pH 7.4) equal to twice the combined weight of the livers. The livers were minced with scissors and homogenized. The homogenate was centrifuged at 10,000 x g (0-4°C) for 20 minutes. The 10,000 x g supernatant was then centrifuged at 100,000 x g for 60 minutes (0-4°C). Following ultracentrifugation, the 100,000 x g supernatant was discarded and the microsomal pellet was resuspended in Tris buffer (0.5 ml/g of liver). The protein content and N-demethylase activity were determined (AW SOP-OP-STX-87, "Determination of Total Protein and N-demethylase Activity"). The microsomes and 100,000 x g supernatant were quickly frozen in liquid nitrogen and stored at -80°C until used.

Incubations

Incubations were prepared using five hepatic subcellular fractions: PB-induced 100,000 x g supernatant,

untreated 10K supernatant, PB-induced microsomes, untreated microsomes, and Arochlor-induced 9K supernatant. The 10K supernatant and microsomal incubations were conducted at a protein concentration of 4 mg/ml. The 2-ml incubates contained 1.0 mM NG or NsG (2 mg/ml DMSO), 5 mM glucose-6-phosphate, 5 mM $MgCl_2$; 1 unit/ml glucose-6-phosphate dehydrogenase, 0.1M pH 7.4 Tris buffer, and the liver fractions. Incubates also contained 0.5 mM NADP whereas the control incubates did not. Additional experiments in which the enzyme system was heat-inactivated were conducted by immersing the subcellular fraction in a 90°C water bath for 10-15 minutes before adding the other cofactors.

In order to simulate the activating conditions for mouse lymphoma studies, the 9K incubations contained, in a total volume of 2 ml: 8 mg of protein, 1.0 mM NsG, 3.375 mg/ml isocitrate, 0.6 mg/ml NADP, and Fischer's medium for leukemic mouse cells. Control incubations did not contain NADP. Control incubates containing only Fischer's medium and NsG were also prepared.

All incubations were performed in a water bath shaker at 37°C. The incubation period for the 10K and microsomal fractions was 20 minutes. The 9K supernatant fractions were incubated for 2 hours. After the desired incubation period, the postincubates were placed in an ice bath and terminated by the addition of 250 μ l of 5% $ZnSO_4$; 2 μ mol of the internal standard (IS) methylnitroguanidine⁴ was then added. The incubates were centrifuged for 5 minutes at 4,000 g, and the resulting supernatant was filtered via Millipore Swinnex filtering system (0.45 μ). The filtrate was analyzed via HPLC.

RESULTS

The HPLC traces obtained from the postincubates were clean with no interfering peaks. Nitroguanidine eluted sharply at 4.9 min, NsG eluted at 4.5 min, and the internal standard eluted at 5.8 min. Table 1 shows the HPLC peak area ratios of NG/IS obtained from incubations of nitroguanidine with untreated and PB-induced 10K supernatant and untreated and PB-induced microsomes. No significant difference in the amount of NG remaining in incubations with the NADPH-generating system and in the corresponding control incubations without NADPH was seen. The same results were obtained when the control incubates were heat-inactivated. The standard deviation values give a rough estimate of the precision of the analysis; in the case of NG, the precision is about 5%.

Table 2 shows the HPLC peak area ratios of NsG/IS obtained from incubations of NsG with untreated and PB-treated 10K supernatant and microsomes as well as with commercially obtained Arochlor-induced 9K supernatant. No significant differences are seen in the amount of NsG present in incubations with or without NADP added; again, the same results were obtained when the enzymes of the control incubates were heat-inactivated. Additionally, NsG was not degraded over a longer incubation period (2 hours at 37°, results not shown). In this case, the precision in the measurement of NsG appears to be in the order of 5-10%.

DISCUSSION

The metabolism of NG and NsG by untreated and phenobarbital-treated rat hepatic 10,000 x g supernatant and microsomal fractions was examined. The disappearance of NG or NsG from these incubations was monitored by HPLC; the results indicate that these compounds do not undergo NADPH-dependent metabolism by rat liver 10,000 x g supernatant or microsomes. This observation is consistent with recent *in vivo* data that indicate NG to be rapidly absorbed through the gut, to enter the bloodstream, and to be quickly excreted unchanged into the urine; it appears that NG and NsG are not metabolized by the liver but are rapidly passed out of the body (2). No metabolism was seen in incubations conducted with hepatic fractions obtained from PB-induced animals, thus demonstrating that NG and NsG are not subject to hepatic enzyme biotransformation.

Nitrosoguanidine was found to have slight mutagenic capabilities in mouse lymphoma assays (1). The metabolism of NsG was examined under conditions similar to that of the mouse lymphoma assay with Fischer's medium for leukemic cells of mice and 9K supernatant to determine if metabolic activity could assist in explaining NsG mutagenicity. No NADPH-dependent metabolism of NsG was seen in this system. This seems to indicate that the mutagenicity observed in the mouse lymphoma assay is not due to a metabolic or chemical degradation product of NsG. However, it should be noted that, due to the imprecision (CV is less than 5%) of the assay in quantitating NsG, small amounts (up to 10 ug) of NsG may be metabolized and yet go undetected under these conditions. Additionally, possible metabolites of NsG are likely to have low molar extinction coefficients so they could be difficult to detect with an UV detector.

REFERENCES

1. Harbell JW. Personal communication.
2. Ho B, Tillotson TA, Sincannon LC, Simboli PB, Korte DW.
The fate of nitroguanidine in the rat. Toxicol Appl
Pharmacol (in preparation).

TABLE 1

The NADPH-Dependent Metabolism of Nitroguanidine by
Untreated and Phenobarbital-Treated (PB) Rat Hepatic
Fractions

Peak Area Ratio of NG/IS \pm SD (n=7)			

		+NADP	-NADP

10K Supernatant			
Untreated	1.10 \pm 0.01	1.12 \pm 0.01	
PB	1.07 \pm 0.04	1.03 \pm 0.06	
Microsomes			
Untreated	1.02 \pm 0.02 ^a	1.03 \pm 0.01 ^b	
PB	0.99 \pm 0.02	0.99 \pm 0.02	

a N = 6

b N = 5

TABLE 2

The NADPH-Dependent Metabolism of Nitrosoguanidine by
 Untreated, Phenobarbital-Treated (PB), and Arochloro-Induced
 Rat Hepatic Fractions

Peak Area Ratio of Nsg/IS \pm SD (n=7)		
	+NADP	-NADP
10K Supernatant		
Untreated	1.00 \pm 0.09	1.00 \pm 0.01
PB	0.86 \pm 0.06	0.86 \pm 0.07
Microsomes		
Untreated	0.89 \pm 0.02	0.99 \pm 0.01
PB	1.33 \pm 0.05	1.37 \pm 0.02
9K Supernatant	0.96 \pm 0.01	0.93 \pm 0.01

N = 5

END

9-87

DTIC